

Electrodynamics in the presence of an axion

C. Corianò *

Institute for Theoretical Physics
and

Department of Physics
State University of New York at Stony Brook
Stony Brook, NY 11794 USA

Mod. Phys. Lett, to appear

Abstract

The low energy limit of an axion field coupled to gauge fields is investigated through the behaviour of the gauge fields propagator in a local vacuum angle background. The local (singular) part of the effective action for the axion field is calculated at one loop level. In the case of a timelike, linearly growing axion field, representing a massive axion, we give an asymptotic expansion of the causal propagator and we solve nonlocally for the first coefficient. We show that, for a generic axionic background, short distance propagation of the gauge field is well defined.

Supported by the NSF under grant NSF 91 08054
and by the Department of Energy under Grant No. DE-FG02-88ER40388

*A. Della Riccia fellow

1 Introduction

One of the most interesting aspects of instanton effects in QCD is related to the strong CP problem and why the vacuum angle is extremely small [1].

Various solution have been proposed in the literature for this problem. In one scheme one can set $\theta = 0$ at tree level and then assume that higher order effects in the magnitude of θ are negligible [2], while a second option is to introduce a massless quark so that θ becomes unobservable. This second possibility is ruled out phenomenologically.

A third alternative is to introduce a dynamical variable $\theta(x)$ in such a way that an effective interaction

$$L_{int} = c\theta(x)F\tilde{F} \quad (1)$$

(c being a constant)

dynamically sets the vacuum angle θ_{QCD} to be zero [3].

This third approach, generally known as the Peccei-Quinn solution, solves the CP problem and has cosmological implications. In this scenario θ is therefore made local, where nonperturbative QCD effects produce a mass for the axion and fix the background value of θ_{QCD} to be zero.

The interaction of an axion field to electromagnetic fields was initially investigated several years ago, after it was shown [4] that in a θ vacuum a t'Hooft-Polyakov magnetic monopole acquires an electric charge proportional to θ and to the magnetic charge g of the monopole,

$$q_\theta = C\theta g \quad (2)$$

with C a constant. In particular the dynamics of monopoles traversing axionic domain walls has since been elucidated [5], [6].

In its full generality, the interaction of the axion with the electromagnetic field is assumed to be given, at low energies, by the action density [5]

$$L = -\frac{1}{4}F^2 + \frac{1}{2}\partial_\mu\theta\partial^\mu\theta - \frac{m^2v^2}{N^2}[1 - \cos(Na/v)] + \alpha F\tilde{F} \quad (3)$$

where $\tilde{F}$ denotes the dual of the field strength and

$$\alpha = \frac{e^2}{32\pi^2 \sin^2\theta_w^0} \left(\theta_{qcd} + \frac{T_\theta}{2\pi} \frac{N\theta(x)}{v} \right) \quad (4)$$

where $\theta(x)$ is the axion field, m is the axion mass, N is the number of axion vacua and v is the vacuum expectation value that breaks the Peccei-Quinn symmetry.

θ_w^0 is the electroweak angle at the grand unification mass scale (*GUM*) and θ_{cd} , the constant is taken at GUM.

$T_\theta = 2\pi$ for QCD, is the period of θ .

We have assumed that there is grand unification of the strong and electroweak interactions [7] and we have used

$$g_s = e^2 / \sin^2(\theta_w^0)$$

at GUM, where g_s is the QCD coupling constant at the same scale.

In this letter we investigate the propagation of high frequency electromagnetic modes in the presence of an axion. We use the light cone expansion of the propagator for the electromagnetic field, and we calculate the local part of the one-loop effective action of the axion field. In the case of a slowly varying θ field, representing a massive axion, we show that the coefficient of the leading singularity of the asymptotic expansion can be explicitly determined in full nonlocal form. This last result allows us to define a causal gauge field propagator in the presence of a massive axion, a result which cannot be obtained from the dispersion relation because it contains spurious poles. As noticed previously in the literature [8], this particular background configuration of the axion field (the massive axion) makes the propagation of the low frequency em modes unstable and a tachionic pole arises. The investigation of ref.[8] was carried out in the particular case of a coupled Maxwell-Chern-Simons theory, a system which shares the same properties of the massive axion model.

We show that, for any background axion field, and in particular for a massive axion model, the em propagator behaves well at short distances. Our results are based on a direct application of the Hadamard theorem [9] for equations of propagation which are diagonal in the highest derivatives. The short distance behaviour of the model is in fact controlled by the highest derivatives of the gaussian operator (which is diagonal), while the instability is generated by the presence of lower derivatives (coupled to an ϵ tensor) in the equations of motion.

The partition function of the model can be written in the simplified form

$$Z = \int [d\theta][dA_\mu] e^{i \int \mathcal{L} + \mathcal{J}\theta + \mathcal{J}'_\mu \mathcal{A}^\mu d^4x} \quad (5)$$

$$\mathcal{L} = -\frac{1}{4}F^2 + \frac{1}{2}\partial_\mu\theta\partial^\mu\theta + \frac{c}{4}\theta(x)F\tilde{F} \quad (6)$$

F denotes the electromagnetic field, $\tilde{F} = \frac{1}{2}\epsilon F$ denotes its dual.

We will work in the Lorentz gauge. The evaluation of the effective action for the θ -field, functionally integrating out the gauge fields is given by

$$\mathcal{L}_{eff} = \frac{1}{2}\partial_\mu\theta\partial^\mu\theta + \frac{i}{2}\text{tr} \ln(\hat{P}) \quad (7)$$

where

$$\hat{P} = g^{\nu\beta}\square - c\partial_\mu\theta\epsilon^{\mu\nu\alpha\beta}\partial_\alpha \quad (8)$$

is the relevant operator induced by the gaussian approximation.

The equations of motion for the gauge fields are:

$$\partial_\mu F^{\mu\nu} = c\partial_\mu\theta\epsilon^{\mu\nu\alpha\beta}\partial_\alpha A_\beta \quad (9)$$

where

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad (10)$$

In the Lorentz gauge we find

$$\square A_\nu - k_{\nu\alpha\beta}\partial^\alpha A^\beta = 0 \quad (11)$$

where we have defined

$$k_{\nu\alpha\beta} = c\partial_\mu\theta\epsilon^{\mu\nu\alpha\beta} \quad (12)$$

The Hadamard expansion [9] for the propagator of the gauge fields in the axionic background can be set directly by introducing the Green's function $\hat{\Delta}$ for eq.(11), since the gaussian operator is diagonal in the highest derivatives

$$\square\hat{\Delta}_{\nu\rho} - k_{\nu\alpha\beta}\partial^\alpha\hat{\Delta}^\beta_\rho = \delta_{\nu\rho}\delta^4(z) \quad (13)$$

where $z = x - y$. The support of the Δ distribution is therefore inside the light-cone and causality is respected.

The standard ansatz is therefore

$$\begin{aligned}\hat{\Delta}_{\nu\rho}(x, y) &= G_{\nu\rho}^{(0)}(x, y)D_F \\ &\quad - \frac{i}{16\pi^2} \ln\left(\frac{z^2}{\mu^2} - i0\right) \sum_{n=0}^{\infty} \left(\frac{z^2}{4}\right)^n \frac{1}{n!} G_{\nu\rho}^{(n+1)}(x, y)\end{aligned}\quad (14)$$

where

$$D_F = \frac{1}{4\pi^2 i(z^2 - i0)} \quad (15)$$

is the Feynman free propagator in coordinate space while μ is a mass parameter introduced in order to keep dimensionless the argument of the logarithm in the expansion.

The recursion relations for the coefficients of the expansion are easily obtained.

By equating to zero the independent singularities of eq. (13) applied to the expansion (14) we get a leading singularity equation for z^{-4}

$$2z^\mu \partial_\mu G_{\nu\rho}^{(0)}(x, y) - z^\alpha k_{\nu\alpha\beta} G_{\beta\rho}^{(0)}(x, y) = 0 \quad (16)$$

a D_F (i.e. z^{-2}) singularity equation:

$$\begin{aligned}\square G_{\nu\rho}^{(0)} + G_{\nu\rho}^{(1)} + z^\mu \partial_\mu G_{\nu\rho}^{(1)} - k_{\nu\alpha\beta} \partial^\alpha G_{\beta\rho}^{(0)} \\ - \frac{z^\alpha}{2} k_{\nu\alpha\beta} G_{\beta\rho}^{(1)\beta} = 0\end{aligned}\quad (17)$$

and a log-equation (for $z^{2n} \ln(\frac{z^2}{\mu^2})$)

$$\begin{aligned}z^\mu \partial_\mu G_{\nu\rho}^{n+2} + (n+2) G_{\nu\rho}^{(n+1)} + \square G_{\nu\rho}^{(n+1)} \\ - \frac{z^\alpha}{2} k_{\nu\alpha\beta} G_{\beta\rho}^{(n+2)\beta} - k_{\nu\alpha\beta} \partial^\alpha G_{\beta\rho}^{(n+1)} = 0\end{aligned}\quad (18)$$

Defining $G^{(-1)} = 0$, $n = -1, 0, 1, \dots$, we summarize all the equations in the form:

$$\begin{aligned}z^\mu \partial_\mu G_{\nu\rho}^{(n+1)} + (n+1) G_{\nu\rho}^{(n+1)} + \square G_{\nu\rho}^{(n)} \\ - \frac{z^\alpha}{2} k_{\nu\alpha\beta} G_{\beta\rho}^{(n+1)\beta} - k_{\nu\alpha\beta} \partial^\alpha G_{\beta\rho}^{(n)} = 0\end{aligned}\quad (19)$$

At this level the axion is a background field and we see, for instance from eq. (16), that it affects the propagation of the leading singularity in a non trivial way (through $k_{\nu\alpha\beta}$).

For a constant θ -angle the equation for the propagator of the leading singularity is trivially given by

$$z^\mu \partial_\mu G_{\nu\rho}^{(0)} = 0 \quad (20)$$

valid on the light cone surface, with the initial condition

$$G_{\nu\rho}^{(0)}(x, x) = \delta_{\nu\rho} \quad (21)$$

The solution is trivially given as the Kronecker delta, and the strength therefore is diagonal over the entire characteristic surface. In this simplified case the equations of motion for the gauge fields are given by

$$\square \square^{-1} = \delta_{\nu\rho} \delta^4(z) \quad (22)$$

where as usual,

$$\square^{-1} = \frac{G_{\nu\rho}^{(0)}}{(z^2 - i0)4\pi^2 i} = \frac{\delta_{\nu\rho}}{(z^2 - i0)4\pi^2 i} \quad (23)$$

At this point we focus our attention on eq (16), which describes the leading behaviour of the dynamics of the electromagnetic field in the presence of a local vacuum angle.

We will preliminarily show that, in the case of a linearly growing axion field, in the timelike case, some components of the leading singularity tensor $G_{\nu\rho}^{(0)}$ are constant on the light cone surface. The reason appears to be quite simple and is valid both in the abelian and in the nonabelian case. In fact, by contraction of both terms of (16) with $\partial_\nu \theta$ and using a symmetry argument (the antisymmetry of k defined in eq (12)) we get the equation

$$\partial_\nu \theta z^\mu \partial_\mu G_{\nu\rho}^{(0)}(x(s), 0) = 0 \quad (24)$$

If we introduce the parametrization

$$x^\mu(s) = sx^\mu, 0 < s < 1; y = 0 \quad (25)$$

for a straight line inside the light cone surface we can rewrite eq (24) in the form

$$\partial_\nu \theta(x) \frac{d}{ds} G^{(0)}_{\nu\rho}(x(s), 0) = 0 \quad (26)$$

Repeating the same procedure a second time one gets the equation

$$x^\nu \frac{d}{ds} G^{(0)}_{\nu\rho}(x(s), 0) = 0 \quad (27)$$

It is simple to show that if the variation of the θ field is linear in a timelike direction, a frame can be found in which it has only a time variation, say

$$\partial_\mu \theta = (a_0, 0, 0, 0) \quad (28)$$

and in particular we get for the timelike components

$$\frac{d}{ds} G^{(0)}_{0\rho} = 0 \quad (29)$$

From the initial condition eq (21) therefore we get

$$G^{(0)}_{0i} = 0 \quad G^{(0)}_{00} = 1 \quad (30)$$

As we emphasized previously, this result remains true even in the more realistic non abelian case, for a linearly growing (timelike) axion field, and is a simple consequence of the antisymmetry involved in the coupling. A simple and physical way to look at this result is to view it as an anisotropy effect induced on the propagation of the gauge fields by the pseudo tensor coupling.

We will show now that the other components of the leading singularity tensor can also be determined by following an approach analogous to the one we used above, in the case in which condition (28) is enforced.

The equation for the leading singularity (eq. 16) can be cast in the form

$$\frac{d}{ds} G^{(0)}_{ij}(x(s), 0) - \frac{ca_0}{2} x^k \epsilon^{0ikl} G^{(0)}_{lj}(x(s), 0) = 0 \quad i, j, k, l = 1, 2, 3 \quad (31)$$

For a given timelike vector x^μ , the previous equation has two first integrals. This can be easily seen through an elementary analogy with the motion

of a classical charged particle under a Lorentz force in a constant magnetic field. We use this analogy to solve it. Define

$$G_{ij}^{(0)}(x(s), 0) = \hat{v}_j; \quad \omega^k = \frac{a_0}{2} x^k \quad (32)$$

where $\hat{\omega}$ is constant at fixed x^μ and rewrite it as

$$\frac{d}{ds} \hat{v}_j = \hat{\omega} \wedge \hat{v}_j \quad (33)$$

and two first integrals of motion along the s-line are

$$\hat{\omega} \cdot \hat{v}_j = \frac{a_0}{2} x^k G_{kj}^{(0)}(x(s), 0) = d_j \quad (34)$$

$$v^2_{\hat{j}} = \sum_i G_{ij}^{(0)}(x(s), 0) G_{ij}^{(0)}(x(s), 0) = l_j \quad (35)$$

Iterating eq.(33) we get

$$\left(\frac{d^2}{ds^2} + \omega^2 \right) \hat{v}_j = \hat{\omega} d_j \quad (36)$$

By using the initial condition eq (21) one easily gets

$$d_j = \frac{ca_0}{2} x^j \quad l_j = 1, \quad j = 1, 2, 3 \quad (37)$$

and finally the solution

$$G_{ij}^{(0)}(x(s), 0) = \left(\delta_{ij} - \frac{x^i x^j}{r^2} \right) \cos\left(\frac{ca_0}{2} rs\right) + \epsilon_{ikj} \frac{x^k}{r} \sin\left(\frac{ca_0}{2} rs\right) + \frac{x^i x^j}{r^2} \quad (38)$$

This expression describes the behavior of the gauge field correlator in its non local form, asymptotically around the light-cone.

Our results can be summarized in the following expressions for the Feynman propagators of electromagnetic fields in the linearly growing timelike θ vacuum

$$< A_i(x) A_j(0) >_\theta =$$

$$\left[\left(\delta_{ij} - \frac{x^i x^j}{r^2} \right) \cos\left(\frac{a_0 r}{2}\right) + \epsilon^{ikj} \frac{x^k}{r} \sin\left(\frac{a_0 r}{2}\right) + \frac{x^i x^j}{r^2} \right] \frac{1}{4\pi^2 i(x^2 - i0)} + \log. terms \quad (39)$$

$$< A_0(x) A_i(0) >_\theta = \frac{\delta_{0i}}{4\pi^2 i(x^2 - i0)} + G_{0i}^{(1)}(x, 0) \log(x^2 - i0) \quad (40)$$

modulo additional logarithmic corrections. This model provides an additional example of a solution for the Green function in an external field in full nonlocal form. The outline of this approach, as is well known, dates back to Schwinger [11] and it dealt with the Dirac propagator in an abelian gauge field. Causality problems have been investigated by Velo and Zwanziger [10] in this framework successfully, and calculations of anomalies are also possible. The massive axion model suffers, however, of an instability at low frequencies, in other words the retarded em propagator develops a tachionic behaviour in its lower modes. This aspect has been considered in very detail in ref. [8] in which the 3+1 dimensional Maxwell-Chern-Simons (MCS) system is discussed. The equations of motion for the propagation of electromagnetic fields in the massive axion background are equivalent, in fact, to the MCS model. An expression of the retarded propagator was also given in it and it was shown that em frequencies smaller than a_0 , the time component of the derivative of the axion field, becomes tachionic [8]. We refer to that paper for further details. In the MCS case the instability is an inevitable effect, since no background field is involved from the beginning, while in the massive axion model is still unclear whether by taking into account in a better way the axion dynamics the instability can be removed.

In its current formulation, in the case of either a spacelike or timelike variation of the local vacuum, the propagation of the gauge fields appears to be consistent in the ultraviolet regime and, therefore, the singular part of the effective action for the axion field, obtained by integrating out the gauge fields in the partition function, can be computed straightforwardly.

In the following we are going to show that the structure of the singularities of the propagator of the gauge field in an arbitrary θ background has a very simple feature.

2 The effective action

In the general case of a variable vacuum angle we can easily calculate the singular (local) part of the effective action for the θ field integrating out the gauge field. Let's define $[G_{\mu\nu}^{(n)}]$ to be the coincidence limit of the coefficients introduced before.

$$[G_{\mu\nu}^n] \equiv G_{\mu\nu}^{(n)}(x, x) \quad (41)$$

The same coefficients also appear in the divergent part of the one loop effective action

$$\Gamma^{(1)}(\theta) = \frac{i}{2} \text{tr} \ln(\hat{P}) \quad (42)$$

Let's take a variation of (42) with respect to the background field θ

$$\delta\Gamma^{(1)}[\theta] = \frac{i}{2} \text{tr} \delta P P^{-1} = \frac{i}{2} < \delta \hat{P}_{\mu\nu} \hat{\Delta}^{\nu\mu}(x, y) \delta^4(z) > \quad (43)$$

where the angular bracket denotes integration over x, y .

We find after some algebra

$$[\delta \hat{P}_{\nu\beta} \hat{\Delta}^{\beta\nu}] = -c < \partial_\mu \delta\theta \epsilon^{\mu\rho\alpha\nu} \partial_\alpha \Delta_{\nu\rho}(x, y) \delta^4(z) > \quad (44)$$

Ultraviolet divergences in the effective action come from the singular behaviour of the propagator at short distances. We need to regularize expressions such as $[\ln(z^2 - i0)]$, $[D_F]$ and $[\partial_\mu D_F]$. This can be done in many ways. The simplest one consists in introducing a short distance cutoff Λ . All the singularities are regulated at the intermediate stage, while we first compute the coincidence limits of the various coefficients. The cutoff is removed by taking the limit $\Lambda \rightarrow \infty$ at the end.

A simple calculation shows that the relevant contribution to (44) is given by

$$[\partial_\alpha \hat{\Delta}_{\nu\rho}] = -\frac{i}{16\pi^2} [\ln(z^2 - i0)] [\partial_\alpha G_{\nu\rho}^{(1)}] + [\partial_\alpha G_{\nu\rho}^{(0)}] [D_F] + [G_{\nu\rho}^{(0)}] [\partial_\alpha D_F] \quad (45)$$

For notational simplicity we omitted to write down in eq. (45) explicitly the cutoff dependence of the singular coincidence limits.

It can be seen immediately that, in this regularization, the contribution from the $[\partial D_F]$ singularity is zero.

By differentiating the singularities equations we find

$$[\partial_m G_{\nu\rho}^{(0)}] = \frac{1}{2} k_{\nu m\rho} \quad (46)$$

$$\begin{aligned} [\partial_m \partial_n G_{\nu\rho}^{(0)}] &= \frac{1}{4} \partial_n k_{\nu m\rho} + \frac{1}{4} \partial_m k_{\nu n\rho} \\ &\quad \frac{1}{8} k_{\nu m}{}^\beta k_{\beta n\rho} + \frac{1}{8} k_{\nu n}{}^\beta k_{\beta m\rho} \end{aligned} \quad (47)$$

The explicit expression for the counterterm is

$$\langle [\delta \hat{P}_{\tau\nu} \hat{\Delta}^{\nu\tau}] \rangle = \int d^4x \delta \partial_\mu \theta \epsilon^{\mu\tau\alpha\nu} \left(\frac{-c}{32\pi^2 i} [\ln(z^2 - i0)] [\partial_\alpha G_{\nu\tau}^{(1)}] + [\partial_\alpha G_{\nu\rho}^{(0)}] [D_F] \right) \quad (48)$$

A straightforward but lengthy calculation gives

$$\Gamma^{(1)}(\theta) = \frac{i}{2} \int d^4x \left(\frac{-c}{32\pi^2 i} [\ln z^2] \left(\frac{1}{4} (Q^2)^2 c^2 - \frac{1}{2} \partial_\mu \theta \square \partial^\mu \theta \right) + \frac{1}{4\pi^2 i} \left[\frac{1}{z^2} \right] \frac{3c}{2} Q^2 \right) \quad (49)$$

where we've defined

$$Q^2 = \partial_\mu \theta \partial^\mu \theta \quad (50)$$

Introducing a pole and a logarithmic renormalization constants Z_1 and Z_2 , we can express the counterterm lagrangean in the form

$$\nabla \mathcal{L}^{count.} = cZ_1 Q^2 + cZ_2 \left(c^2 (Q^2)^2 - 2 \partial_\mu \theta \square \partial^\mu \theta \right) \quad (51)$$

3 Conclusions

We have analyzed the dynamics of a coupled axion-gauge field model in the low energy limit and we've investigated in detail the problem of the propagation of high frequency em modes in an axion background. In particular we have seen that, in the case of a timelike varying axion background, the leading singularity of the Hadamard expansion of the gauge fields correlator can be determined in non local form. By applying Hadamard's theorem on the structure of the singularities of diagonal hyperbolic operators, we have shown that the instability of the Maxwell-axion system is confined to the

propagation of low em frequencies, reobtaining a result first considered in ref. [8]. Our results are valid for the propagation of em modes in any axionic background. We have also shown that the local part of the effective action of the axion can be computed in a simple form, a result which gives to the effective model physical consistency.

Acknowledgments

I acknowledge Profs. A.S. Goldhaber, H. Yamagishi, B. W. Lindquist J.J.M. Verbaarschot and G. Sterman. I warmly thank Prof. D. Zwanziger and Dr. F. Bastianelli for clarifying discussions and for helpful suggestions. I finally thank the Theory Group at Univ. of Lecce, Italy for their kind hospitality, Susan Marie Trapasso and Rinat Kedem for their generous help.

References

- [1] G. t'Hooft Phys. Rev.D14, 3432 (1976); C. G. Callan, R. F. Dashen and D. J. Gross, Phys Lett.63B, 334 (1976); R. Jackiw and C. Rebbi, Phys. Rev. Lett. 37, 172 (1976).
- [2] R. Crewther, P. Di Vecchia, G. Veneziano and E. Witten Phys. Lett.B88, 123 (1979); V. Baluni, Phys. Rev. D19, 2227.
- [3] R.D. Peccei and H.R. Quinn, Phys. Rev. Lett.38, 1440 (1977) and Phys. Rev. D16, 1791 (1977); S. Weinberg, Phys Rev. Lett.40, 223 (1978).
- [4] E. Witten, Phys.Lett. 86B, 283 (1979).
- [5] P. Sikivie, Phys. Lett. 137B 353 (1984).
- [6] M.C. Huang and P. Sikivie, Phys. Rev. D32, 1560 (1985); P. Simic, Nucl. Phys. B272, 365 (1986).
- [7] H. Georgi and S.L. Glashow, Phys. Rev. Lett.48, 1867 (1982).
- [8] S. M. Carroll, G. B. Field and R. Jackiw, Phys. Rev. D41, 1231 (1990).
- [9] J. Hadamard, in "Lectures on Cauchy's Problem in Linear Partial Differential Equations" (New Haven:Yale Univ. Press) (1913). B. De Witt, in "Dynamical Theory of Groups and Fields" (Gordon and Breach, London) (1965).
- [10] G. Velo and D. Zwanziger in "Invariant wave equations" A. S. Wightman ed. (Springer Verlag Lecture Notes 73) (1977).
- [11] J. Schwinger, Phys. Rev 93 664 (1951).